

Kidney Failure (ESRD)

Disease Report

REPORT CATEGORY —



Sample Client

Report date: 15 January 2026

Table of Contents

03 How this works

- 04 Impact
- 05 Evidence
- 06 Some things to keep in mind

07 Introduction

08 Your genetics

09 Your recommendations

33 Next Steps

- 33 Your Lab Results

DISCLAIMER

This report does not diagnose this or any other health conditions. Please talk to a healthcare professional if this condition runs in your family, you think you might have this condition, or you have any concerns about your results.

Personal information

NAME

Sample Client

SEX AT BIRTH


Male

HEIGHT

5ft 5" 165cm


WEIGHT

137lb 62kg



How this works

Our Wellness Reports analyze how your DNA influences your health.
We then use this analysis to give you personalized risk estimates and recommendations.




Similarly, our Trait Reports look at how your DNA influences your traits.



Your DNA is like an instruction manual — it contains a lot of information.
You can think of it as a blueprint for your body.

Genetic variants are parts of DNA that differ from person to person. Some can make you more vulnerable to certain health issues, while others may influence traits such as eye color.



We use artificial intelligence and machine learning to analyze all this information.
We then summarize your results as a risk score or display it on a gauge.

In total, we analyze up to 83 million genetic variants.

When we give a risk score, the risk icon tells you if you are at a higher or lower risk compared to other people:



Genotype color info:

AA

You don't have any risk alleles

AA

You have 1 risk allele

AA


You have 2 risk alleles

Your risk is also displayed as a percentile. This will tell you how your risks compare to our sample population. The lower your percentile number, the lower your risk. The "50th percentile" would be an average risk.

Similarly, the gauge tells you your relative risk score compared to our sample population, or it indicates a specific trait or haplotype you are more likely to have based on your genetic variants.

When applicable, we also list top evidence-based recommendations that may help lower your risk. The focus is on recommendations that may be of benefit to you, based on your genetics.

Our recommendations come in four categories: lifestyle, diet, supplements and drugs.
The following icons tell you which category a recommendation falls into:



Our team of scientists also ranks each recommendation.
We rank based on impact and the strength of evidence in the medical literature.

Impact shows how strongly a recommendation will affect your health in a certain area.
Evidence is how much scientific support there is for the recommendation. Rankings are from 1 to 5 (low to high):



Impact

Impact scores range from 1-5. These scores reflect how much of an effect each recommendation can have. An impact score of 5 predicts the biggest effect.

When a recommendation affects something we can measure, we use those measurements to assign the impact score. For example, a recommendation that decreases cholesterol by 20% will have a higher impact score than one that decreases it by 5%.

Some recommendations affect things that we cannot directly measure, like stress or mood. For these, the impact score is based on how well they work relative to other recommendations and standard treatments. The best ones get the highest scores.

If there is a lot of research that shows a recommendation works especially well for your genotype, the impact score gets increased.

Recommendation Evidence

- 5 / 5
- Recommendations that are considered effective and generally recommended by experts and medical bodies.
-
- 4 / 5
- Recommendations that are considered likely effective and that have multiple independent meta-analyses and a great many studies supporting them.
-
- 3 / 5
- Recommendations that are considered possibly effective and have many studies supporting them
-
- 2 / 5
- Recommendations that have insufficient evidence, with two or several clinical trials supporting them, or many studies but with ambiguous results.
-
- 1 / 5
- Recommendations that have insufficient evidence, with a single clinical trial, or with many studies most of which didn't find support for the recommendation.
-
- 0 / 5
- No evidence in humans.

Genotype-specific Evidence

- High-quality
- Direct evidence that a recommendation helps more in people with your gene variant (many clinical trials, a few large clinical trials, or a meta-analysis).
-
- Medium-quality
- Direct evidence that a recommendation helps more in people with your gene variant (a few clinical trials or one large clinical trial).
-
- Low-quality
- Direct evidence that a recommendation helps more in people with your gene variant (a single clinical trial or more trials with inconsistent results).
-
- Indirect
- A recommendation may help more in people with your gene variant because it targets a specific gene or protein affected by your variant (e.g., MTHFR, dopamine).
-
- In theory
- A recommendation may help more in people with your gene variant because it targets a specific mechanism affected by your variant (e.g., inflammation, oxidative stress).

Some things to keep in mind:

- Genetics doesn’t play a considerable role in a condition or a trait.
- There is not enough research available to estimate a genetic predisposition.
- There are technical limitations to estimating or presenting a genetic predisposition.
- The topic is sensitive, and a genetic predisposition should only be estimated and presented by a healthcare professional.

Introduction

Kidney failure is a condition that occurs when the kidneys can no longer filter waste products from the blood.

Signs and symptoms of kidney failure include:

- Fatigue
- Weakness
- Nausea and vomiting
- Loss of appetite
- Shortness of breath
- Swelling
- High blood pressure
- Anemia
- Bone disease

Risk Factors and Treatment

Conditions that may cause kidney failure include:

- Diabetes
- High blood pressure
- Glomerulonephritis (an inflammation of the kidneys)
- Polycystic kidney disease
- Kidney stones
- Urinary tract infections

Managing these underlying conditions is crucial to slow the progression of kidney damage. Medications may be prescribed to control blood pressure, reduce proteinuria, and manage other symptoms. Dialysis, either hemodialysis or peritoneal dialysis, may be necessary to filter waste products and excess fluids from the blood in cases of advanced kidney failure. In some instances, a kidney transplant may be considered as a long-term solution.

Alongside medical interventions, lifestyle changes like maintaining a balanced diet, limiting salt and protein intake, staying hydrated, and avoiding tobacco and excessive alcohol consumption can significantly improve kidney function and overall health. Regular monitoring by healthcare professionals is essential to adjust treatment plans as needed and optimize outcomes.



MORE LIKELY

More likely to have kidney failure based on 15,508 genetic variants we looked at

Your Recommendations

Your recommendations are prioritized according to the likelihood of it having an impact for you based on your genetics, along with the amount of scientific evidence supporting the recommendation.

You’ll likely find common healthy recommendations at the top of the list because they are often the most impactful and most researched.

DOSAGE		DOSAGE	
1	Avoid Dioxin	2	Aerobic Exercise (Cardio)1 hour
3	Cordyceps500 mg	4	Avoid Mycotoxin
5	N-acetylcysteine (NAC)1200 mg	6	Walking30 minutes
7	Exercise At Least One Hour a Day1 hour	8	Mindfulness Meditation30 minutes
9	B Vitamins	10	Limit Salt Intake
11	Avoid Refined Sugar	12	Stress Management Therapy1 hour
13	Yoga30 minutes	14	Limit Meat Intake
15	DASH Diet	16	Limit Protein Intake
17	Maintain Optimal Vitamin D Levels1000 iu	18	Good Oral Hygiene2 minutes
19	Sleep for 7+ Hours	20	Avoid Magnesium Supplements
21	Probiotics30 billion CFU	22	Avoid Sugary Foods & Drinks
23	Avoid Processed High Phosphate Foods	24	Tai Chi1 hour
25	Coenzyme Q10 (CoQ10)100 mg	26	Reduce Oxalate Intake
27	L-Carnitine1 g	28	Omega-3 (Fish Oil)2000 mg

1



Avoid Dioxin

IMPACT4 / 5

EVIDENCE2 / 5

How to implement

Reduce consumption of animal fats, since dioxins accumulate in fat tissue. Choose lean cuts of meat, and opt for organic or pasture-raised when possible to minimize exposure. Additionally, avoid burning trash that contains plastic, treated wood, or chlorinated chemicals to prevent dioxin release into the environment.

Description

Avoiding exposure to dioxin, a highly toxic environmental pollutant, can help protect against serious health risks, including cancer and hormonal disruptions. Reducing contact with dioxin-contaminated sources such as certain foods and industrial emissions is essential for long-term health.

How it helps

A large study associated exposure to the dioxins 1,2,3,6,7,8-hexachlorodibenzo-p-dioxin, PCB 126, PCB 169, and PCB 156 with a 7-fold increased risk of nephropathy. Women exposed to dioxins were at even higher (17-fold) risk [\[R\]](#).

2



Aerobic Exercise (Cardio)

IMPACT

3 / 5

EVIDENCE

3 / 5

How to implement

Engage in at least 150 minutes of moderate-intensity aerobic exercise or 75 minutes of vigorous-intensity activity each week. Distribute this time over at least 3 days per week, avoiding consecutive days of vigorous exercise to allow for recovery.

TYPICAL STARTING DOSE

1 hour

Description

Engaging in regular aerobic exercise, such as running, swimming, or cycling, offers numerous health benefits, including improved cardiovascular fitness, weight management, and mood enhancement. It supports overall physical and mental well-being while reducing the risk of chronic diseases.

Cardio, short for cardiovascular exercise, is any type of physical activity that temporarily increases your heart rate. Examples include **running, cycling, swimming, and brisk walking**.

Regular cardio exercise has many benefits for your overall health. It can help lower your risk of heart disease and diabetes, support weight loss, and improve your mood and energy levels. To get the most out of cardio, try to do it for at **least 30 minutes, 3-5 times a week**.

Interval training is a type of cardio that combines periods of high-intensity training with brief rest periods.

How it helps

Experts agree that 30-60 minutes of moderate exercise most days of the week may support kidney health. Increased physical activity is linked to a lower risk of kidney failure [\[R, R, R\]](#).

Exercise may help by [\[R, R, R, R, R, R, R\]](#):

- Improving physical fitness
- Boosting kidney function
- Reducing blood pressure

3



Cordyceps

IMPACT

2 / 5

EVIDENCE

3 / 5

How to implement

Take 500-1000 mg of Cordyceps extract in supplement form once daily, preferably with a meal to enhance absorption. This dosage is commonly recommended for general wellness and should be continued on a long-term basis, as effects accumulate over time.

TYPICAL STARTING DOSE

500 mg

Description

Cordyceps is a type of medicinal mushroom known for its potential to boost endurance and enhance respiratory function. It's used in traditional medicine and dietary supplements to support physical performance and overall vitality.

[Cordyceps](#) are a group of fungi (mushrooms) that grow on insects. Some species, like *O. sinensis* and *C. militaris*, are used in supplements [\[R\]](#).

O. sinensis grows on a moth caterpillar. Extracts and teas made from the fungus-caterpillar combination are often used in traditional Chinese medicine. They are thought to boost sex drive, reduce fatigue, and support kidney health [\[R\]](#), [\[R\]](#), [\[R\]](#).

How it helps

Two meta-analyses (the largest one with 9 studies) concluded that supplementation with cordyceps (3-6 g/day for 6-24 months) **improves graft function, anemia, leucopenia, and liver function while reducing acute rejection episodes, lung infection, serum albumin and uric acid levels, hyperlipidemia, hyperglycemia, proteinuria, CNI nephrotoxicity, and hepatotoxicity** [\[R\]](#), [\[R\]](#).

4



Avoid Mycotoxin

IMPACT

2 / 5

EVIDENCE

2 / 5

How to implement

To avoid mycotoxins, store grains and nuts in a dry, cool place, check for mold on foods before eating, and avoid consuming products with visible mold. Prefer buying whole foods over processed ones, and if possible, opt for foods tested for mycotoxin contamination. Discard any food that smells musty or shows signs of spoilage.

Description

Avoiding mycotoxin exposure, which can occur in contaminated foods, is crucial to prevent potential health risks such as liver damage and cancer. Being cautious about food quality and storage conditions can help minimize the risk of mycotoxin ingestion.

How it helps

Exposure to ochratoxin A has been associated with endemic nephropathy [\[R\]](#), [\[R\]](#), [\[R\]](#).

5



N-acetylcysteine (NAC)

IMPACT1 / 5

EVIDENCE2 / 5

How to implement

Take 600 mg of N-Acetylcysteine (NAC) supplement daily with water. It can be taken at any time of the day, but try to take it at the same time each day for best results.

TYPICAL STARTING DOSE
1200 mg

Description

NAC is a supplement that contains a form of the amino acid cysteine, a protein building block that your body uses to make the antioxidant glutathione. It is used for its potential antioxidant properties and its ability to support lung, gut, and mental health.

[N-acetylcysteine](#) (NAC) is converted to cysteine in the body. Cysteine is a protein building block (amino acid) that helps make the antioxidant glutathione [\[R\]](#).

People take NAC to potentially support [\[R\]](#), [\[R\]](#):

- Mental health
- Ovarian health and pregnancy outcomes
- Lung health
- Gut health

How it helps

Patients undergoing cardiac surgery, especially those with pre-existing kidney issues, are at risk of developing acute kidney injury. NAC has been studied in this setting for its potential to enhance blood flow and protect kidney function. However, evidence regarding its effectiveness remains inconclusive.

The typical dosage of NAC varies based on the clinical scenario. Oral NAC is often used, with common regimens including 600 mg to 1200 mg twice daily. Intravenous formulations are also available, particularly in acute settings like post-heart surgery.

While NAC is generally well-tolerated, it's important to consider potential side effects like nausea, vomiting, and rare allergic reactions. Additionally, its effectiveness in kidney protection is not definitively established, and it should not replace other preventive measures like adequate hydration.

References: [R](#), [R](#), [R](#), [R](#), [R](#), [R](#), [R](#)

6

Walking

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Incorporate at least 30 minutes of brisk walking into your daily routine, aiming for a minimum of five days a week. This can be done in one continuous session or broken into shorter periods, such as three 10-minute walks throughout the day.

TYPICAL STARTING DOSE

30 minutes

Description

Walking is a low-impact form of exercise that can contribute to cardiovascular fitness, weight management, and improved overall health. It is used to support physical activity goals, enhance mood, and promote better cardiovascular health.

How it helps

Regular walking can improve cardiovascular health, control weight, and reduce stress, all of which help manage symptoms and complications of kidney failure.

7

Exercise At Least One Hour a Day

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Dedicate a minimum of 60 minutes to moderate-intensity activities such as brisk walking, swimming, or cycling. Do this most days of the week, aiming for at least 5 days to optimize benefits.

TYPICAL STARTING DOSE

1 hour

Description

[Exercise](#) can do wonders for your health. It can help you lose weight, improve your heart health, boost your mood, and more [\[R\]](#).

There are many ways you can be active. You can walk, run, swim, dance, or play team sports. **Everything counts, and it's never too late to start!**

Try to get at least **60 minutes of moderate physical activity on most days**, including walking. **Getting a mix of cardio (min 150 min/day) and strength training (min 2 times a week) may be optimal.**

How it helps

Regular exercise can help improve cardiovascular health and control blood pressure, which are important for patients with kidney failure.

8



Mindfulness Meditation

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Practice mindfulness meditation for 10-20 minutes daily. Find a quiet, comfortable place to sit or lie down, then focus on your breath, observing thoughts and sensations without judgment. Consistency is key, so try to incorporate it into your daily routine, perhaps in the morning or before bed.

TYPICAL STARTING DOSE

30 minutes

Description

Mindfulness meditation involves focused attention on the breath or a specific point of focus to cultivate mindfulness and reduce stress. Regular meditation practice is associated with improved mental clarity, emotional regulation, and overall well-being.

How it helps

Mindfulness meditation can reduce stress and improve mental health, which can be beneficial for managing symptoms and overall well-being in kidney failure patients.

9



B Vitamins

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Take a B vitamin complex supplement once daily, preferably with your morning meal to enhance absorption. The supplement should include a range of B vitamins such as B1 (thiamine), B2 (riboflavin), B3 (niacin), B5 (pantothenic acid), B6 (pyridoxine), B7 (biotin), B9 (folate), and B12 (cobalamin). Ensure consistency by taking it at the same time each day.

Description

B vitamins help the body function properly. Two examples are [vitamin B9 \(folate\)](#) and [vitamin B12](#). They play essential roles in [\[R, R, R, R\]](#):

- DNA formation
- Metabolism
- Energy production

You can get these vitamins from [\[R, R, R, R, R\]](#):

- Animal products
- Green leafy vegetables
- Citrus fruits
- Fortified foods
- Supplements

Adults should be getting **400 micrograms (mcg) of folate** and **2.4 mcg of vitamin B12** every day [\[R, R\]](#).

Folate deficiency is rare but can happen in people that don’t eat enough fruits and vegetables. Alcoholics and lactating mothers may also be at an increased risk [\[R\]](#).

Vitamin B12 deficiency often takes years to develop, as the body is able to store large amounts in the liver. Groups that may be at an increased risk include [\[R, R\]](#):

- Vegans
- Pregnant women
- Older people
- People with gut issues

How it helps

B Vitamins are essential for maintaining overall kidney health. They can help support metabolic processes, reduce inflammation, and improve energy levels in patients with kidney disease.

10



Limit Salt Intake

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Reduce your daily salt intake to less than 2,300 milligrams (mg), equivalent to about one teaspoon of table salt. This includes all salt consumed, whether added during cooking, at the table, or present in processed and ready-to-eat foods.

Description

Limiting salt intake involves lowering the amount of sodium in the diet, which can help lower blood pressure and reduce the risk of heart disease. It often involves avoiding processed and other high-sodium foods such as cured meats, soups and sauces, and various pickled foods.

Your body needs a certain amount of sodium to function. For example, it helps support your blood pressure. This sodium usually comes from [salt](#). However, eating too much salt is linked to an increased risk of stroke and heart disease [\[R\]](#), [\[R\]](#).

Foods higher in salt include [\[R\]](#), [\[R\]](#), [\[R\]](#):

- Prepared soups, broths, gravies, and sauces
- Fermented or pickled foods (such as pickles, soy sauce, olives, and [capers](#))
- Cured meats and sausages (such as bacon, salt pork, and salami)
- Cheese (such as Roquefort and Parmesan)

Try to limit your daily intake to 1 teaspoon of salt (2,400 mg of sodium) [\[R\]](#).

How it helps

Reducing salt intake can help control blood pressure, which is crucial for people with kidney failure. High blood pressure can cause further damage to weakened kidneys.

11



Avoid Refined Sugar

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Eliminate foods and beverages high in refined sugars, such as sodas, candies, pastries, and white bread, from your diet. Replace them with natural sugars found in fruits and whole grains. Implement this change immediately and maintain it indefinitely for long-term health benefits.

Description

Reducing the consumption of refined sugar and high-sugar foods can help prevent dental problems, manage blood sugar levels, and support weight management.

How it helps

Refined sugar can lead to spikes in blood sugar levels, which can worsen kidney function and overall health in ESRD patients.

12



Stress Management Therapy

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Engage in stress management therapy sessions, such as cognitive-behavioral therapy (CBT), for at least 1 hour per week over a course of 8 to 12 weeks. Techniques can include mindfulness, deep breathing exercises, and identifying stressors to develop coping strategies.

TYPICAL STARTING DOSE

1 hour

Description

Stress management therapy refers to various techniques and approaches aimed at reducing and coping with stress. It can improve mental and physical well-being by helping individuals better manage the effects of stress on their health.

How it helps

Managing stress can lower blood pressure and improve heart health, which is beneficial for individuals with kidney failure.

13



Yoga

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Practice yoga for at least 20 to 30 minutes a day, most days of the week. Choose a style that matches your fitness level and goals, and consider attending a class or using online resources to guide your practice.

TYPICAL STARTING DOSE

30 minutes

Description

Yoga is a mind-body practice that combines physical postures, breathing exercises, and meditation. It enhances flexibility, strength, and mental well-being and is used for stress reduction, relaxation, and overall health improvement.

[Yoga](#) combines breathing, stretching, and relaxation techniques. Practicing yoga may help [R](#), [R](#), [R](#):

- Reduce [stress](#)
- Improve fitness
- Lower blood pressure and heart rate
- Manage pain

How it helps

Yoga can help improve flexibility, reduce stress, and enhance overall quality of life for people with kidney failure.

14



Limit Meat Intake

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Substitute red and processed meats with plant-based proteins, such as beans, lentils, and tofu, for at least two meals per week. Gradually increase the number of meat-free meals by incorporating more fruits, vegetables, and whole grains into your diet daily.

Description

Limiting meat intake involves consuming fewer animal-based products, often as part of a plant-based diet. This dietary choice can provide health benefits such as lower cholesterol levels, reduced risk of certain chronic diseases, and a decreased environmental footprint.

Proteins, fats, and carbs are macronutrients. You need these nutrients in large (‘macro’) amounts. They provide your body with the energy it needs to function properly [\[R\]](#).

You should choose high-quality foods that give you the “best” version of each macronutrient. For example, too much saturated fat and protein from certain animal sources may increase the risk of heart disease and infertility [\[R, R, R\]](#).

Consider replacing red meat with other high-protein foods, such as [\[R, R\]](#):

- Eggs
- Fish
- Soy
- Beans

How it helps

Reducing meat intake can decrease the workload on kidneys by limiting the amount of protein waste the kidneys need to filter. This helps manage waste buildup in the blood.

15



DASH Diet

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Adopt a dietary pattern that emphasizes fruits, vegetables, whole grains, lean proteins, and low-fat dairy, while reducing sodium, red meat, sweets, and sugary beverages. Aim for 4-5 servings of both fruits and vegetables per day, 6-8 servings of grains (with at least half being whole grains), and 2-3 servings of low-fat dairy. This diet should be followed daily to manage blood pressure effectively.

Description

The DASH (Dietary Approaches to Stop Hypertension) diet is a dietary plan designed to reduce high blood pressure. It emphasizes fruits, vegetables, lean proteins, and whole grains while limiting sodium intake.

Health experts developed the DASH diet to reduce blood pressure. This diet also helps support weight control while reducing inflammation and cholesterol [\[R\]](#).

It **limits salt, sweets, and saturated fat**, while promoting the intake of [\[R\]](#):

- Fruits (4-5 servings/day)
- Vegetables (4-5 servings/day)
- Whole grains (6-8 servings/day)
- Low-fat dairy (2-3 servings/day)
- Nuts, seeds, and legumes (4-5 servings/week)
- Fish and poultry (up to 6 ounces/day)

The DASH diet focuses on fresh, minimally processed food rich in nutrients. Magnesium, calcium, and potassium-rich foods are particularly important due to their beneficial effects on the heart and blood vessels [\[R\]](#).

Although similar to the [Mediterranean diet](#), DASH is lower in sodium (<2,300 mg a day) and omega-3 fatty acids. On the other hand, it’s higher in red meat and dairy [\[R\]](#).

How it helps

The DASH diet, which is low in sodium and rich in fruits and vegetables, can help reduce blood pressure. For those with kidney failure, managing blood pressure is essential to slowing down the progression of kidney damage.

16



Limit Protein Intake

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Limit your daily protein intake to 0.8 grams per kilogram of your body weight. For example, if you weigh 70 kilograms (about 154 pounds), aim to consume no more than 56 grams of protein per day. Adjust this goal based on your activity level and health status, reducing intake further if advised by a healthcare professional.

Description

Proteins, fats, and carbs are macronutrients. You need these nutrients in large (‘macro’) amounts. They provide your body with the energy it needs to function properly [\[R\]](#).

You should choose high-quality foods that give you the “best” version of each macronutrient. For example, too much protein from certain animal sources may increase the risk of bone, kidney, and metabolic issues in some people [\[R\]](#), [\[R\]](#), [\[R\]](#).

How it helps

Excessive protein can produce more waste products that the failing kidneys will struggle to remove, making ESRD worse.

17



Maintain Optimal Vitamin D Levels

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Check your vitamin D levels, they should ideally be in the 30-66 ng/mL range. If your levels are lower than that, take a vitamin D supplement, 1000-4000 IU daily, to reach an optimal range.

TYPICAL STARTING DOSE

1000 iu

Description

Vitamin D, often referred to as the "sunshine vitamin," can be synthesized by the skin when exposed to sunlight, as well as being found in fish, eggs, and fortified milk. It helps regulate calcium absorption, promoting strong bones and a healthy immune system. Vitamin D deficiency can lead to conditions like rickets in children and osteoporosis in adults.

Your body needs [vitamin D](#) for strong bones. Vitamin D also plays a role in [\[R\]](#):

- Mood
- Immunity
- Heart health
- Blood sugar control

[Sunlight](#) is our main source of vitamin D. Experts recommend getting at least **5-15 minutes of midday sun, 2-3 times per week**. People with darker skin and those living at high latitudes may need longer periods of sun exposure [\[R, R\]](#).

Foods like fish, eggs, and fortified milk provide small amounts of vitamin D. **People lacking vitamin D should consider taking a supplement** [\[R\]](#).

How it helps

Vitamin D supplementation helps with calcium and phosphorus balance, which is often disrupted in kidney failure, reducing the risk of bone disease.

18



Good Oral Hygiene

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Brush your teeth at least twice a day for two minutes each time using fluoride toothpaste. Floss daily to remove plaque from places your toothbrush can't reach and consider using mouthwash. Visit your dentist at least twice a year for professional cleanings and checkups.

TYPICAL STARTING DOSE

2 minutes

Description

Good oral hygiene, which includes regular brushing, flossing, and dental check-ups, is crucial for preventing tooth decay, gum disease, and maintaining overall oral health. A healthy mouth contributes to overall well-being and can prevent dental issues that may impact systemic health.

Oral hygiene refers to practices that keep the mouth healthy. Good habits include [R](#), [R](#):

- Gently brushing the teeth and tongue for two minutes, at least twice a day
- Flossing daily
- Using a toothpaste with fluoride and a soft toothbrush
- Replacing a worn toothbrush at least every 3 months

To help maintain good oral hygiene, it is also important to regularly see dental professionals. They can help prevent and address issues with your teeth and gums.

How it helps

Poor oral health is linked to chronic kidney disease. Maintaining good oral hygiene helps reduce bacteria that can cause infections, potentially reducing the risk of kidney damage.

19



Sleep for 7+ Hours

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Ensure you allocate enough time in your schedule to achieve a minimum of 7 hours of sleep each night. This might involve going to bed earlier or adjusting your evening routine to promote relaxation and make it easier to fall asleep.

Description

Optimizing sleep involves adopting healthy sleep habits and creating a sleep-conducive environment to ensure restorative and sufficient sleep duration. It supports cognitive function, mood stability, and overall physical health. Most experts recommend getting **at least 7 hours of good-quality sleep each night**.

[Sleep](#) supports your body and mind [\[R, R\]](#). More precisely, sleep helps:

- Support brain health [\[R, R\]](#)
- Maintain a healthy weight and appetite [\[R, R, R\]](#)
- Regulate blood pressure [\[R, R\]](#)
- Balance blood sugar [\[R, R\]](#)

Ways to sleep better include [\[R\]](#):

- Reducing your bright light exposure (screen time) in the evenings
- Sticking to a regular sleep schedule
- Avoiding hunger or large meals before bed
- Avoiding nicotine, caffeine, and alcohol before bed
- Maintaining a sleep area that’s cool, dark, and quiet

How it helps

Adequate sleep helps regulate crucial bodily functions, which can reduce the burden on kidneys and improve overall health and well-being.

20



Avoid Magnesium Supplements

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Do not purchase or consume magnesium supplements, including tablets, capsules, or powders. If you are currently taking magnesium supplements, consider discontinuing their use, especially if not medically advised.

Description

Magnesium supplements may not be necessary if you're eating a balanced diet, as it's found in many foods. Overuse can lead to stomach issues, irregular heartbeat, and even cardiac arrest. It's best to get your magnesium naturally through diet, unless a doctor recommends otherwise.

How it helps

In kidney failure, the kidneys are not able to effectively filter and remove excess magnesium from the body. Taking magnesium supplements can lead to harmful accumulation, worsening the condition. It's crucial for patients with kidney failure to avoid extra magnesium to prevent its buildup.

0 / 5

22



Avoid Sugary Foods & Drinks

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

To avoid sugary foods, eliminate or significantly reduce consumption of foods and beverages high in added sugars such as sodas, candies, baked goods, and sugary cereals from your diet. Instead, opt for natural sugar sources like fruits. Aim to do this daily for ongoing health benefits.

Description

High-sugar foods like baked goods, sweets, and sugary drinks, can spike your blood sugar levels. Consuming a lot of these types of foods can contribute to health issues like diabetes, obesity, insomnia, and heart disease.

High-sugar foods and refined carbs have a high glycemic index (GI). This means they tend to spike your blood sugar levels. They include [R](#), [R](#), [R](#):

- Sugary drinks
- Baked goods
- Sweets
- White bread
- White rice
- Pasta

Eating a lot of sugary foods can contribute to:

- Diabetes [R](#), [R](#), [R](#)
- Weight gain and obesity [R](#), [R](#)
- Insomnia [R](#)
- Heart disease [R](#)

You may also want to avoid processed sugars and sugary drinks. They may have a role in increasing IL-1B [R](#), [R](#).

How it helps

Sugary foods can lead to increased blood sugar levels, which can strain the kidneys further and potentially contribute to worsening kidney function in people with kidney failure.

23



Avoid Processed High Phosphate Foods

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

To follow this recommendation, remove or significantly reduce processed foods such as sodas, packaged snacks, processed meats, and canned foods from your daily diet. Instead, opt for fresh fruits, vegetables, whole grains, and lean proteins. It's helpful to read nutrition labels, aiming to keep your daily phosphate intake below 700 mg if possible.

Description

Processed foods often have high levels of phosphate. These include fast foods and soda. Extra phosphate can make bones brittle and damage your blood vessels over time. So, it's healthier to choose fresh foods over processed ones.

How it helps

Eating foods high in phosphates can be harmful for those with kidney failure because damaged kidneys can't remove phosphorus from the blood effectively. This can lead to high levels of phosphorus in the blood, causing bone and cardiovascular issues.

24



Tai Chi

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Practice Tai Chi for 30 to 60 minutes at least twice a week. Choose a quiet, spacious area and follow along with a qualified instructor, either in person at a class or through an online video tutorial, to ensure proper technique and maximum benefit.

TYPICAL STARTING DOSE

1 hour

Description

Tai Chi is a traditional Chinese mind-body practice involving slow, flowing movements and deep breathing. It is known for its potential to reduce stress, improve balance, and enhance overall physical and mental well-being.

Tai chi involves gentle movements and breathing to strengthen and relax the mind and body. Practicing tai chi may help [\[R\]](#), [\[R\]](#), [\[R\]](#):

- Manage pain
- Improve fitness
- Increase well-being
- Improve sleep and mood

How it helps

Tai Chi can improve cardiovascular health, reduce stress, and enhance physical function, which are important for managing kidney failure.

25



Coenzyme Q10 (CoQ10)

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Take a 100 mg Coenzyme Q10 (CoQ10) supplement once daily with a meal that contains fat for better absorption.

TYPICAL STARTING DOSE

100 mg

Description

Coenzyme Q10 (CoQ10) is a naturally occurring antioxidant that plays a crucial role in cellular energy production. It is often taken as a supplement to support heart health, improve energy levels, and provide antioxidant protection, especially for individuals with certain medical conditions or as they age.

[Coenzyme Q10](#) (CoQ10) is a compound that helps enzymes work better. By doing so, CoQ10 helps improve [R](#), [R](#), [R](#), [R](#):

- Energy levels
- Antioxidant protection
- Heart health
- Muscle strength
- Blood sugar control

The amount of CoQ10 made by your body decreases as you get older. Luckily, you can also get it from food or supplements. Good sources of CoQ10 include [R](#), [R](#):

- Organ meats
- Fatty fish
- Whole grains

How it helps

Coenzyme Q10 (CoQ10) has antioxidant properties that can help protect the kidneys from further damage and improve kidney function. It can also help in reducing oxidative stress and inflammation.

26



Reduce Oxalate Intake

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Avoid high-oxalate foods such as spinach, beets, potatoes, and nuts. Instead, choose low-oxalate options like bananas, cabbage, and cucumbers. Aim to keep your daily oxalate intake under 100mg.

Description

Reducing oxalate intake means consuming foods and beverages with lower oxalate content, which can help prevent kidney stone formation and support kidney health. Oxalates are naturally occurring compounds found in various foods like spinach, chocolate, and nuts, and managing their intake is important for those prone to kidney stone formation.

Oxalate (*oxalic acid*) is a natural substance found in many foods, such as [\[R\]](#), [\[R\]](#), [\[R\]](#):

- **Spinach**
- Tea
- Rhubarb
- Chocolate
- Beets
- Nuts

Adults typically get 150-250 mg of oxalate every day. A high intake of oxalate may reduce calcium absorption and increase the risk of kidney stones [\[R\]](#), [\[R\]](#), [\[R\]](#).

How it helps

Reducing oxalate intake is important for individuals with kidney failure because oxalates can contribute to kidney stone formation, worsening kidney function. Eating fewer oxalate-rich foods helps reduce the risk of kidney stones.

27



L-Carnitine

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Take 500 mg of L-carnitine supplement daily with a glass of water, preferably with a meal to enhance absorption.

TYPICAL STARTING DOSE

1 g

Description

L-carnitine is an amino acid-like compound that plays a role in energy metabolism and helps prevent toxic substances from building up in cells. It is often used in dietary supplements for its potential to support muscle recovery, reduce fatigue, and enhance athletic performance.

[L-carnitine](#) is a compound that helps you burn fat. It also prevents toxic substances from building up in cells [\[R\]](#).

Your body can usually make enough carnitine to meet its needs. You can also get it from **meat and dairy products** [\[R\]](#).

People use L-carnitine for [\[R\]](#), [\[R\]](#), [\[R\]](#), [\[R\]](#), [\[R\]](#), [\[R\]](#):

- Heart problems
- Overactive thyroid
- Fertility problems
- Blood sugar control
- Weight control

How it helps

L-Carnitine can help with muscle weakness and fatigue that are common in ESRD patients. Its supplementation has been shown to improve energy levels by aiding in the transport of fatty acids into the mitochondria for energy production.

How to implement

TYPICAL STARTING DOSE

2000 mg

Description

[Omega-3 fatty acids](#) are some of the healthiest fats we can eat. They help lower inflammation and protect the heart, brain, and eyes. Our bodies produce less omega-3s than we need for optimal health, so it's important to get enough through food or supplements [[R](#), [R](#), [R](#)].

There are three major types of omega-3s: ALA, EPA, and DHA [R, R].

Fatty fish are rich in EPA and DHA. They include [\[R\]](#):

- Salmon
- Tuna
- Herring
- Sardines

For optimal protection, try to get at least **two servings of fatty fish per week**. Fish oil supplements are available for those who don't eat fish regularly [R].

How it helps

Omega-3 fatty acids can help reduce inflammation and improve heart health, which is often compromised in ESRD. Studies also suggest they might help lower blood pressure and triglycerides, providing cardiovascular benefits.

Next Steps

Remember, your genes only tell one important part of your health story!


Now that you've seen your DNA-based results for this health topic, let's take a look at other contributing factors.

Your lab results

Your lab results are impacted by the combined effect of your genes, environment and lifestyle.

Lab tests will give you the best picture of your current health status, while your genes provide insight into your health predispositions and which recommendations are best for you.

	Protein, Urine		Negative Positive	9 Nov 2025
	Phosphorus	2 mg/dL	12.53.54.5	9 Nov 2025
	WBC	6.92 x10^3/mm3	246.51140	9 Nov 2025
	eGFR	71 mL/min/1.73 m2	306090120	29 Aug 2025
	Zinc	17.12 umol/L	7.659.1812.2419.89	9 Nov 2025
	BUN/Creatinine Ratio	13 x100%	101520	9 Nov 2025
	Vitamin D, 25-Hydroxy, Total	58.59 ng/mL	10203075150	9 Nov 2025
	Albumin	4.3 g/dL	3.44.45.5	9 Nov 2025
	Creatinine	1.08 mg/dL	0.60.71.11.2	9 Nov 2025

	Potassium, Serum	5.2 mmol/L		9 Nov 2025
	Albumin/Creatinine Ratio, Random Urine	37 mg/g crt		9 Nov 2025
	Albumin, Urine	87.6 ug/mL		9 Nov 2025