

Spinal Canal Narrowing

DNA Health Report

REPORT CATEGORY —



BONE HEALTH

Sample Client

Report date: 15 January 2026

Powered by

 omicse Edge

Table of Contents

03 How this works

- 04 Impact
- 05 Evidence
- 06 Some things to keep in mind

07 Introduction

08 Your genetics

09 Your recommendations

DISCLAIMER

This report does not diagnose this or any other health conditions. Please talk to a healthcare professional if this condition runs in your family, you think you might have this condition, or you have any concerns about your results.

Personal information

NAME

Sample Client

SEX AT BIRTH

Male

HEIGHT

5ft 5" 165cm

WEIGHT

137lb 62kg



How this works

Our Wellness Reports analyze how your DNA influences your health.
We then use this analysis to give you personalized risk estimates and recommendations.



Similarly, our Trait Reports look at how your DNA influences your traits.



Your DNA is like an instruction manual — it contains a lot of information.
You can think of it as a blueprint for your body.

Genetic variants are parts of DNA that differ from person to person. Some can make you more vulnerable to certain health issues, while others may influence traits such as eye color.



We use artificial intelligence and machine learning to analyze all this information.
We then summarize your results as a risk score or display it on a gauge.

In total, we analyze up to 83 million genetic variants.

When we give a risk score, the risk icon tells you if you are at a higher or lower risk compared to other people:



Genotype color info:

AA

You don't have any risk alleles

AA

You have 1 risk allele

AA

You have 2 risk alleles

Your risk is also displayed as a percentile. This will tell you how your risks compare to our sample population. The lower your percentile number, the lower your risk. The "50th percentile" would be an average risk.

Similarly, the gauge tells you your relative risk score compared to our sample population, or it indicates a specific trait or haplotype you are more likely to have based on your genetic variants.

When applicable, we also list top evidence-based recommendations that may help lower your risk. The focus is on recommendations that may be of benefit to you, based on your genetics.

Our recommendations come in four categories: lifestyle, diet, supplements and drugs.
The following icons tell you which category a recommendation falls into:



Our team of scientists also ranks each recommendation.
We rank based on impact and the strength of evidence in the medical literature.

Impact shows how strongly a recommendation will affect your health in a certain area.
Evidence is how much scientific support there is for the recommendation. Rankings are from 1 to 5 (low to high):



Impact

Impact scores range from 1-5. These scores reflect how much of an effect each recommendation can have. An impact score of 5 predicts the biggest effect.

When a recommendation affects something we can measure, we use those measurements to assign the impact score. For example, a recommendation that decreases cholesterol by 20% will have a higher impact score than one that decreases it by 5%.

Some recommendations affect things that we cannot directly measure, like stress or mood. For these, the impact score is based on how well they work relative to other recommendations and standard treatments. The best ones get the highest scores.

If there is a lot of research that shows a recommendation works especially well for your genotype, the impact score gets increased.

Recommendation Evidence

- 5 / 5
- Recommendations that are considered effective and generally recommended by experts and medical bodies.
-
- 4 / 5
- Recommendations that are considered likely effective and that have multiple independent meta-analyses and a great many studies supporting them.
-
- 3 / 5
- Recommendations that are considered possibly effective and have many studies supporting them
-
- 2 / 5
- Recommendations that have insufficient evidence, with two or several clinical trials supporting them, or many studies but with ambiguous results.
-
- 1 / 5
- Recommendations that have insufficient evidence, with a single clinical trial, or with many studies most of which didn't find support for the recommendation.
-
- 0 / 5
- No evidence in humans.

Genotype-specific Evidence

- High-quality
- Direct evidence that a recommendation helps more in people with your gene variant (many clinical trials, a few large clinical trials, or a meta-analysis).
- Medium-quality
- Direct evidence that a recommendation helps more in people with your gene variant (a few clinical trials or one large clinical trial).
- Low-quality
- Direct evidence that a recommendation helps more in people with your gene variant (a single clinical trial or more trials with inconsistent results).
- Indirect
- A recommendation may help more in people with your gene variant because it targets a specific gene or protein affected by your variant (e.g., MTHFR, dopamine).
- In theory
- A recommendation may help more in people with your gene variant because it targets a specific mechanism affected by your variant (e.g., inflammation, oxidative stress).

Some things to keep in mind:

- Genetics doesn’t play a considerable role in a condition or a trait.
- There is not enough research available to estimate a genetic predisposition.
- There are technical limitations to estimating or presenting a genetic predisposition.
- The topic is sensitive, and a genetic predisposition should only be estimated and presented by a healthcare professional.

Introduction

Spinal stenosis is a condition characterized by the narrowing of spaces within the spine, which can put pressure on the nerves that travel through the spine. The narrowing can occur in any part of the spine but is most commonly found in the lumbar (lower back) and cervical (neck) areas. This condition often arises from wear-and-tear changes in the spine related to osteoarthritis.

In severe cases, this compression can lead to a significant decrease in quality of life due to pain, numbness, or even weakness in the limbs. Walking and standing for long periods often exacerbates the symptoms, which can lead to a progressive loss of mobility.

Diagnosis and Treatment

The symptoms of spinal stenosis can vary widely, from mild discomfort to severe impairment. Some individuals may experience back pain, tingling, weakness, or numbness that can radiate from the back into the buttocks and legs, commonly referred to as sciatica when it affects the sciatic nerve.

Diagnosis typically involves physical examination, assessment of medical history, and imaging studies such as MRI or CT scans to view the degree of canal narrowing and nerve compression. Treatment options range from conservative measures, like physical therapy and medications, to interventional procedures such as epidural steroid injections, and in advanced cases, surgery to relieve the pressure on the nerves might be considered.



TYPICAL LIKELIHOOD

Typical likelihood of spinal stenosis based on 1,001,035 genetic variants we looked at



Your top variants that most likely impact your genetic predisposition:

GENE	SNP	GENOTYPE
IPP	rs2356581	AA
DYNC111	rs2051706	GG
TRIM32	rs1040851	CC
PDE3A	rs7298820	AT
SMAD3	rs28613758	CT
SLITRK6	rs2183749	AC
ELMOD2	rs184579809	TT
HEY1	rs143958409	CC
AGMO	rs7808567	CC
GFPT1	rs6728752	AA
CALHM6	rs2637678	CC

The number of "risk" variants in this table doesn't necessarily reflect your overall result.

Your Recommendations

Your recommendations are prioritized according to the likelihood of it having an impact for you based on your genetics, along with the amount of scientific evidence supporting the recommendation.

You’ll likely find common healthy recommendations at the top of the list because they are often the most impactful and most researched.

DOSAGE		DOSAGE	
1	Avoid DHEA Supplements	2	Dietary Omega-3 Fatty Acids
3	Stretching15 minutes	4	Dietary Magnesium
5	Aquatic Exercise1 hour	6	Physical Therapy30 minutes
7	Avoid High-Impact Exercise	8	Topical Capsaicin

1



Avoid DHEA Supplements

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Do not purchase or consume any supplements labeled as DHEA (dehydroepiandrosterone). If you currently have DHEA supplements, stop using them immediately and do not resume.

Description

If you're taking DHEA supplements, reduce the dosage, or stop taking them until your levels go back to normal.

How it helps

Genetically higher testosterone levels may increase the risk of spinal stenosis.

The study utilized Mendelian randomization to assess the phenome-wide effects of free testosterone in males from the UK Biobank. Lifelong increased testosterone had mixed effects, including benefits on bone density and adverse effects on prostate cancer and spinal stenosis [\[R\]](#).

2



Dietary Omega-3 Fatty Acids

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Incorporate foods high in omega-3 fatty acids into your diet daily. This includes eating fish such as salmon, mackerel, and sardines at least twice a week. Alternatively, include a tablespoon of flaxseed oil or chia seeds in your daily diet.

Description

[Omega-3 fatty acids](#) are some of the healthiest fats we can eat. They help lower inflammation and protect the heart, brain, and eyes. Our bodies produce less omega-3s than we need for optimal health, so it's important to get enough through food or supplements [\[R\]](#), [\[R\]](#), [\[R\]](#).

There are three major types of omega-3s: ALA, EPA, and DHA. **EPA and DHA are vital for mental and heart health** [\[R\]](#), [\[R\]](#), [\[R\]](#).

Fatty fish are rich in EPA and DHA. They include [\[R\]](#):

- Salmon
- Tuna
- Herring
- Sardines

For optimal health, try to get at least **two servings of fatty fish per week**. Fish oil supplements are available for those who don't eat fish regularly [\[R\]](#).

How it helps

Omega-3 fatty acids, found in fish and certain plant oils, have anti-inflammatory properties that can be beneficial for managing symptoms of spinal stenosis. They help reduce inflammation in the body, including around the spinal cord, potentially alleviating pain and discomfort associated with spinal stenosis.

3



Stretching

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Incorporate stretching exercises into your daily routine, dedicating at least 10-15 minutes each day. Focus on major muscle groups such as the neck, shoulders, chest, back, hips, and legs. For best results, stretch both before and after other physical activities, holding each stretch for 15-30 seconds without bouncing.

TYPICAL STARTING DOSE

15 minutes

Description

Stretching involves gently elongating the muscles to improve flexibility, reduce muscle tension, and enhance joint mobility. Regular stretching can help prevent injuries, promote better posture, and increase overall physical comfort.

Stretching is a form of physical exercise that involves moving a muscle or group of muscles to their maximum range of motion. The most common types are **dynamic** (involving movement) and **static** stretching.

Stretching can help **improve flexibility, range of motion, and posture**. It may also relieve stress, reduce muscle soreness and tension, increase blood flow, and reduce the risk of injuries.

To get the most out of stretching, try to do it regularly, breathe deeply, and ease into the stretches.

How it helps

Regular stretching helps improve flexibility and range of motion in the spine and surrounding muscles. This can alleviate the pressure on the nerves affected by spinal stenosis, reducing symptoms like pain and numbness.

4



Dietary Magnesium

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Increase your intake of magnesium-rich foods such as leafy green vegetables, nuts, seeds, and whole grains. Aim to include these foods in your diet daily, following the recommended dietary allowance of 320 mg per day for women and 420 mg per day for men.

Description

Magnesium is a vital mineral involved in over 300 biochemical reactions in the body. It supports various functions, including muscle and nerve function, bone health, and blood sugar regulation.

[Magnesium](#) is an essential mineral. Your body needs it for [\[R\]](#), [\[R\]](#), [\[R\]](#):

- Muscle, nerve, and bone function
- Blood sugar and blood pressure control
- DNA and protein production
- Strong immunity

Women need **310-320 mg** of magnesium per day, while men need **400-420 mg** [\[R\]](#).

Foods rich in magnesium include **nuts, seeds, and leafy greens**. Magnesium is also available as a supplement [\[R\]](#).

How it helps

Magnesium plays a key role in maintaining muscle function and bone health. For those with spinal stenosis, adequate magnesium intake can help in managing symptoms by supporting the health of the spinal column and potentially reducing inflammation that contributes to stenosis.

5



Aquatic Exercise

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Participate in aquatic exercise sessions, such as swimming or water aerobics, for 60 minutes, 3 to 5 times per week. Ensure the exercise intensity is moderate, allowing you to talk but not sing during the activity. Consistency over time is key, so aim to incorporate this into your weekly routine for at least 3 to 6 months to observe benefits.

TYPICAL STARTING DOSE

1 hour

Description

Aquatic exercise involves physical activity performed in water, such as swimming, water aerobics, or aquatic therapy. It is gentle on the joints, making it suitable for individuals with mobility issues, and can help improve cardiovascular fitness and muscle strength.

Aquatic exercise is any type of low-impact physical activity performed in a pool. In this type of cardio exercise, water helps the body float, which reduces gravity and makes the practice more comfortable and tolerable [\[R\]](#), [\[R\]](#).

Compared to land-based exercise, aquatic exercise may be better by [\[R\]](#):

- Reducing the stress and impact on joints
- Lowering fracture risk

Hence, it is usually **recommended for the elderly**. Aquatic exercise is also suitable for people who don't know how to swim [\[R\]](#).

How it helps

Exercising in water provides a gentle resistance and supports the body, reducing the stress on the spine and joints. For people with spinal stenosis, aquatic exercises can improve physical function and decrease pain.

6



Physical Therapy

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Attend physical therapy sessions 2-3 times per week for a duration of 4-6 weeks, depending on your specific condition and the advice of your healthcare provider. Each session typically lasts about 30-60 minutes, where a licensed therapist will guide you through targeted exercises, stretches, and possibly other treatments like electrical stimulation or ultrasound therapy.

TYPICAL STARTING DOSE

30 minutes

Description

Physical therapy is a therapeutic practice focused on optimizing physical function and mobility through specialized exercises, manual techniques, and therapeutic modalities. It helps individuals recover from injuries, manage chronic conditions, and improve overall physical well-being by enhancing strength, flexibility, and pain management.

Physical therapy (physiotherapy) helps people regain or maintain their ability to move [\[R\]](#).

Physical therapy can involve [\[R\]](#):

- Joint or muscle exercises
- Corrective movements
- Massage
- Education and advice

People mainly use physical therapy to help with [\[R\]](#), [\[R\]](#):

- Pain and injury
- Stroke recovery
- Chronic health conditions
- Headaches

Mirror therapy uses a mirror placed between the arms or legs. The image of a moving arm or leg gives the illusion of normal movement in the affected one. This therapy stimulates different brain regions and aims to improve mobility [\[R\]](#).

Constraint-induced movement therapy is another type of physical therapy. It consists of restraining the healthy leg or arm to increase the use of the affected one [\[R\]](#).

How it helps

Physical therapy offers a tailored exercise program to strengthen the muscles supporting the spine, improving balance and mobility. This can help manage the symptoms of spinal stenosis and prevent further degeneration.

7



Avoid High-Impact Exercise

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Opt for low-impact activities such as swimming, cycling, or walking instead of running or jumping exercises. Engage in these alternative exercises for at least 150 minutes per week, spread out over several days to ensure rest periods between sessions.

Description

For individuals with certain medical conditions or joint issues, avoiding high-impact exercise can help prevent injuries and joint strain. Choosing lower-impact activities like swimming or cycling can provide similar cardiovascular benefits while minimizing the risk of overuse injuries.

How it helps

High-impact exercises can exacerbate the symptoms of spinal stenosis by increasing the pressure on the spine. Avoiding such activities helps in managing pain and prevents further damage to the spinal structures.

8



Topical Capsaicin

IMPACT

0 / 5

EVIDENCE

0 / 5

How to implement

Apply a capsaicin cream or gel to the affected area 3 to 4 times daily. Make sure the skin is clean and intact before application. Avoid contact with eyes, mouth, and other sensitive areas. Use consistently for several weeks to see improvement.

Description

Capsaicin is a compound found in chili peppers and is used topically in creams and patches for its potential to alleviate pain related to conditions like arthritis, neuropathy, and muscle soreness.

[Capsaicin](#) is the compound that makes chili peppers spicy [\[R\]](#).

People use topical capsaicin to relieve pain and oral capsaicin to aid weight loss [\[R\]](#), [\[R\]](#).

In the short term, capsaicin causes pain. However, **if you’re exposed to a lot of capsaicin over a long period of time, it can reduce your pain sensitivity** [\[R\]](#), [\[R\]](#).

***Please note:** Capsaicin may irritate the skin. Consult your doctor if the irritation is severe or lasts longer than 4 weeks. If capsaicin gets in your eyes, flush them with water immediately and consult your doctor* [\[R\]](#), [\[R\]](#), [\[R\]](#).

How it helps

Capsaicin cream, derived from chili peppers, can reduce pain by depleting the nerve cells of substance P, a chemical component carrying pain messages to the brain. Its analgesic effect is beneficial for managing symptoms in conditions like spinal stenosis, where nerve pain is common.